

Physiotherapy Effects on Pain while Walking in Patients with Knee Osteoarthritis		Healthcare
		Keywords: knee osteoarthritis, gait, Kinesio Tape, rehabilitation, numerical pain rating scale - NRS, 10 meter walking test.

Klejda Tani	University Aldent
Kiri Zallari	Faculty of Technical Medical Sciences, UMT
Irena Kola	Faculty of Technical Medical Sciences, UMT

Abstract

Knee osteoarthritis is a chronic degenerative disease. The main symptom of patients with knee osteoarthritis is a degenerative and mechanics type of pain. Pain is very noticeable while walking in rugged terrain, during ascent and descent of stairs, when changing from sitting in standing position as well as staying in the sitting position for a long time. Many studies have shown that the strength of the quadriceps femoris muscle can affect gait, by improving or weakening it. Kinesio Tape is a physiotherapeutic technique, which reduces pain and increases muscular strength by irritating the skin receptors. The aim of this article was to verify if Kinesio Tape reduces pain while walking, in patients with knee osteoarthritis. 74 patients with primary knee osteoarthritis, aged 50-73 years, participated in this study. We observed the change of pain, while walking for 10 meters at normal speed for each patient, before, a day after the application and three days after the application of Kinesio Tape on quadriceps femoris muscle, with the help of numerical pain rating scale - NRS. Our results indicated that there was a significant decrease of the pain while walking for 10 meters.

1. Introduction

Osteoarthritis is a widespread, slowly developing disease, with a high prevalence increasing with age. The most common large joints involved in the disease are the knees, where the disease is particularly disabling because of difficulty in rising from chair, climbing stairs, kneeling, standing and walking. These limitations are partly due to muscle weakness, especially quadriceps muscle (1–4). It has been suggested that functional ability is also affected by poor proprioception (5 –11).

The primary complaints of patients suffering from osteoarthritis are pain, stiffness, instability and loss of function. In addition to this impaired muscle function is frequently observed in patients with osteoarthritis of the knee. It was found that 80% of patients with knee osteoarthritis reported problems related to muscle function, strength, endurance and coordination (18).

Hassan et al. (12) observed that knee joint pain and quadriceps strength were significant predictors of increased postural sway. Women who reported widespread pain had a 60% greater risk of falling compared to women with no or mild pain (13).

Additional factors associated with disability in persons with knee osteoarthritis include increasing age, obesity, female gender, co morbidity and quadriceps muscle weakness (20, 21). Pain and muscle strength may particularly influence postural sway (12). Impairments in knee joint proprioception have been mentioned by multiple authors (12, 15, 16). These deficits, in combination with the ageing process, may culminate in greater impairments of balance in this patient population, compared with their age-matched and healthy counterparts (13).

Kinesio Tape is the original and authentic elastic therapeutic tape. According to Kenzo Kase (17,29), Kinesio Tape is a definitive rehabilitative taping technique that is designed to facilitate the body’s natural healing process while providing support and stability to muscles and joints without restricting the body’s range of motion as well as providing extended soft tissue manipulation to prolong the benefits of manual therapy administered within the clinical setting. These proposed mechanisms may include: 1. correcting muscle function by strengthening weakened muscles, 2. improving circulation of blood and lymph by eliminating tissue fluid or bleeding beneath the skin by moving the muscle, 3. decreasing pain through neurological suppression, and 4.

repositioning subluxed joints by relieving abnormal muscle tension, helping to return the function of fascia and muscle. Chen et al. (22), mentioned in their studies that Kinesio Tape can lift the skin to increase space between skin and muscle, reducing and localized pressure, promoting circulation and lymphatic drainage. This theoretically reduces pain, swelling, and muscle spasm. Chronic pain can be improved via the sensory stimulation of other types of nerve fibers. In these circumstances, Kinesio Tape may be effective for pain that persists after an injury has healed or for pain that is above and beyond the injury severity

To our knowledge, no study has directly investigated the effect of Kinesio Tape on pain intensity during walking in patients with knee osteoarthritis. The aim of this study was verify if the application of Kinesio Tape on quadriceps muscle reduces pain while walking for 10 meters at normal speed, in patients with knee osteoarthritis before the application of Kinesio Tape, a day after the application of Kinesio Tape and three days after the application of Kinesio Tape on quadriceps femoris muscle.

2. Patients and Methods

The subjects (n=74), aged 50-73years (mean age 61.5), were consecutive out-patients with a clinical diagnosis of primary knee osteoarthritis made by a rheumatologist. The main criterion for the selection of the subjects in this study was the diagnosis of knee osteoarthritis by X-ray. Criteria for excluding subjects in the study were other musculoskeletal diseases, total knee replacement, significant hip or spinal arthritis, neurological diseases and diseases that affect balance and coordination. All of the subjects signed a written consent to participate in the study voluntarily.

Kinesio Tape was applied with a tonus regulation technique also called muscle technique on quadriceps femoris muscle. We measured the tape length in maximal stretched position of the tissue. The application was done with the patient in this maximal stretched position. The tape was applied without stretch following the course of the muscle borders from one insertion to the opposite one.


Fig1. Apply of Kinesio Tape on quadriceps femoris muscle.



We observed the change of pain, while walking for 10 meters at normal speed for each patient, before, a day after the application and three days after the application of Kinesio Tape on quadriceps femoris muscle, with the help of numerical pain rating scale - NRS. The worse knee, as selected by the patient was the “index” knee. Pain was assessed by numerical pain rating scale (NRS), by instructing the patient to choose a number from 0 to 10 that best describes their current pain. 0 would mean “no pain” and 10 would mean “worst possible pain” (19).

In the current study the alternative hypothesis that Kinesio Tape is effective to decrease the pain while walking on a distance of 10 meters in patients with knee osteoarthritis was tested.

Fig2. Numerical pain rating scale used in this study.



3. Results

Seventy four consecutive out-patients with a clinical diagnosis of primary knee osteoarthritis participated in this study, mean age of the participants was 61.5 (range 50-73 years), 67% of whom were female. The worse knee, as selected by the patient was the “index” knee.

We observed the change of pain, while walking for 10 meters at normal speed for each patient before the application of Kinesio Tape, a day after the application of Kinesio Tape and three days after the application of Kinesio Tape on quadriceps femoris muscle, with the help of numerical pain rating scale (NRS), where 0 would mean “no pain” and 10 would mean “worst possible pain” (19).

In Table 1 is shown the number of patients who chose the pain score while walking for 10 meters on normal speed before applying the Kinesio Tape. It is shown that 21 of 74 patients (28.4%) chose score 5, 15 of 74 patients (20.3%) chose score 6, 17 of 74 patients (22.9%) chose score 7 and 21 of 74 patients (28.4%) chose score 8 on the numerical pain rating scale.

Table 1. Numerical pain rating scale (NRS) scores before applying Kinesio Tape (KT)

NRS	Number of patients before KT	%
0		
1		
2		
3		
4		
5	21	28,4
6	15	20,3
7	17	22,9
8	21	28,4
9		
10		

In Table 2 is shown the number of patients who chose the pain score while walking for 10meters on normal speed one day after applying the Kinesio Tape. It is shown that 10 of 74 patients (13.5%) chose score 4, 26 of 74 patients (35.1%) chose score 5, 29 of 74 patients (39.3%) chose score 6, 5 of 74 patients (6.7%) chose score 7 and 4 of 74 patients (5.4%) chose score 8 on the numerical pain rating scale.

Table 2. Numerical pain rating scale (NRS) scores scores one day after applying Kinesio Tape (KT)

NRS	Number of patients one day after KT	%
0		
1		
2		
3		
4	10	13,5
5	26	35,1
6	29	39,3
7	5	6,7
8	4	5,4
9		
10		

In Table 3 is shown the number of patients who chose the pain score while walking for 10meters on normal speed three days after applying the Kinesio Tape. It is shown that 15 of 74 patients (20.3%) chose score 2, 26 of 74 patients (35.1%) chose score 3, and 33 of 74 patients (44.6%) chose score 4 on the numerical pain rating scale.

Table 3. Numerical pain rating scale (NRS) scores three days after applying Kinesio Tape (KT)

NRS	Number of patients three days after KT	%
0		
1		
2	15	20,3
3	26	35,1
4	33	44,6
5		
6		
7		
8		
9		
10		

Table 4. Numerical pain rating scale (NRS) scores before applying Kinesio Tape (KT), one day and three days after applying KT.

NRS	Number of patients before KT	%	Number of patients one day after KT	%	Number of patients three days after KT	%
0						
1						
2					15	20,3
3					26	35,1
4			10	13,5	33	44,6
5	21	28,4	26	35,1		
6	15	20,3	29	39,3		
7	17	22,9	5	6,7		
8	21	28,4	4	5,4		
9						
10						

4. Discussion

Lack of information about the impact of elastic therapeutic tape in pain relief in this diagnosis led us to carry out this research. Our objective was to determine whether the application of Kinesio Tape on quadriceps muscle in patients with knee osteoarthritis will lead to a pain relief while walking a 10 meter distance on a normal speed.

The results of this study showed no significant difference in pain intensity during walking one day after applying Kinesio Tape on quadriceps femoris muscle. However a significant decrease in pain intensity was shown during walking three days after applying the Kinesio Tape. Similar findings have been reported elsewhere. Kaya et al. (23) studied 55 patients with shoulder impingement syndrome treated by Kinesio Tape or local modalities and found that although immediate effect of Kinesio Tape is greater than the local modalities, Kinesio Tape was similarly effective at the second week of the treatment.

The results of the study conducted by Miller and Osmotherly (24) provided evidence for a short-term role for taping as an adjunct to routine physiotherapy program in different treatments. They found that Kinesio Tape has main effect on the early stage of treatment and that there was not a significant Kinesio Tape effect after several weeks. The immediate results and improvements following the Kinesio Tape are also reflected in the work of researchers who found significant improvements immediately following Kinesio Tape compared with placebo taping in patients with other musculoskeletal disorders such as patellofemoral pain syndrome (25,26) or whiplash-associated disorders (27,28).

Kase et al. (17,29) and Thelen et al. (30) however, recommend at least three daily action of elastic therapeutic tape. Kase et al. mentions that three days after the application of Kinesio Tape can occur soft tissue changes, improvement of muscle function, increase of blood circulation and lymphatic drainage. Thelen et al. found that after three days of Kinesio Tape application, was shown a significant decrease of the functional shoulder joint pain and increase of movement.

Limitations in this study was the sample size, with a greater sample size we could get more relevant results. In this study the effect of Kinesio Tape in pain relief in knee osteoarthritis was not assessed. Further studies are needed to investigate the effect of Kinesio Tape in pain relief on knee osteoarthritis.

5. Conclusion

There seems to be a significant decrease of pain three days after applying Kinesio Tape on quadriceps femoris muscle. However, no significant improvement was shown a day after the application. Kinesio Tape can be used in patients with knee osteoarthritis, especially when pain relief is a short term goal of the treatment. More clinical research is needed to investigate the effect of Kinesio Tape in pain relief of knee osteoarthritis.

References

1. Van Baar ME, Dekker J, Lemmens JA, Oostendorp RA, Bijlsma JW. (1998). Pain and disability in patients with osteoarthritis of hip or knee: the relationships with articular, kinesiological, and psychological characteristics. *J Rheumatol* ;25:125–33.
2. Franssen M, McConnell S, Bell M.(2002). Therapeutic exercise for people with osteoarthritis of the hip or knee: a systematic review. *J Rheumatol* ;29:1737–45.
3. Steultjens MP, Dekker J, van Baar ME, Oostendorp RA, Bijlsma JW. (2001). Muscle strength, pain and disability patients with osteoarthritis. *Clin Rehabil* ;15:331–41.
4. Slemenda C, Brandt KD, Heilman DK, Mazucca S, Braunstein EM, Katz BP, et al. (1997). Quadriceps weakness and osteoarthritis of the knee. *Ann Intern Med* ;127:97–104
5. Sharma L, Cahue S, Song J, Hayes K, Pai YC, Dunlop D. (2003). Physical functioning over three years in knee osteoarthritis: role of psychosocial, local mechanical, and neuromuscular factors. *Arthritis Rheum*; 48:3359–70.
6. Sharma L. (2003) Proprioception in osteoarthritis. In: Brandt KD, Doherty M, Lohmander LS, editors. *Osteoarthritis*. (2nd ed). Oxford: Oxford University Press;. p. 172–7.
7. Sharma L, Pai YC, Holtkamp K, Zev Rymer W. (1997). Is knee proprioception worse in the arthritic knee versus the unaffected knee in unilateral knee osteoarthritis? *Arthritis Rheum*; 40:1518–25.
8. Bennell KL, Hinman RS, Metcalf BR, Crossley KM, Buchbinder R, Smith M, et al. (2003). Relationship of knee joint proprioception to pain and disability in individuals with knee osteoarthritis. *J Orthop Res*;21:792–7.
9. Hurley MV, Scott DL, Rees J, Newham DJ. (1997). Sensorimotor changes and functional performance in patients with knee osteoarthritis. *Ann Rheum Dis*;56:641–8.
10. Pai Y, Zev Rymer W, Chang RW, Sharma L. (1997). Effect of age and osteoarthritis on knee proprioception. *Arthritis Rheum*; 40:2260–5.
11. Marks R. (1994). Correlation between knee position sense measurements and disease severity in persons with osteoarthritis. *Rev Rheum Engl Ed*;61:365–72..
12. Hassan B. S., Mockett S., Doherty M. (2001) Static postural sway, proprioception and maximal voluntary quadriceps contraction in patients with knee osteoarthritis and normal control subjects. *Ann. Rheum. Dis.* 60: 612–618
13. Mandeville D., Osternig L. R., Chou L. S. (2008) The effect of total knee replacement surgery on gait stability. *Gait Posture.* 27: 103–109
14. Hinman R. S., Bennell K. L., Metcalf B. R., Crossley K. M. (2002) Balance impairments in individuals with symptomatic knee osteoarthritis: a comparison with matched controls using clinical tests. *Rheumatology (Oxford).* 41: 1388–1394
15. Hurley M. V., Scott D. L., Rees J., Newman D. J. (1997) Sensorimotor changes and functional performance in patients with knee osteoarthritis. *Ann. Rheum. Dis.* 56: 641–648
16. Tarigan T. J., Kasjmir Y. I., Atmakusuma D., Lydia A., Bashiruddin J., Kusumawijaya K., Prihartono J. (2009) The degree of radiographic abnormalities and postural instability in patients with knee osteoarthritis. *Acta Med. Indones.* 41: 15–19
17. Kase, K., Wallis, J., Kase, T.(2003): *Clinical therapeutic applications of the kinesio taping method*. Tokyo: Ken Ikai Co. Ltd.,. 348 p. ISBN 978-1-528725-68-2.

18. Henrik Rogind, MD, Birgitte Bibow-Nielsen, MD, Bodil Jensen, PT, Hans C. Moller, MD, Hans Frimodt-Møller, MD, Henning Bliddal, MD. (1998). The Effects of a Physical Training Program on Patients With Osteoarthritis of the Knees. *Arch Phys Med Rehabil* ;79:1421-1427.
19. McCaffery, M., Beebe, A., et al. (1989). *Pain: Clinical manual for nursing practice*, Mosby St. Louis, MO.
20. McAlindon TE, Cooper C, Kirwan JR, Dieppe PA. (1993). Determinants of disability in osteoarthritis of the knee. *Ann Rheum Dis*;52:258-62.
21. Dekker J, Tola P, Aufdemkampe G, Winckers M. (1993). Negative affect, pain and disability in osteoarthritis patients: the mediating role of muscle weakness. *Behav Res Ther*;31:203-6.
22. Chen PI, Hong WH, Lin CH, Chen WC.(2008). Biomechanics effects of kinesiotaping for persons with patellofemoral pain syndrome during stair climbing. *Biomed.*;21:395-397.
23. Kaya E, Zinnuroglu M, Tugcu I. (2011). Kinesio taping compared to physical therapy modalities for the treatment of shoulder impingement syndrome. *Clin Rheumatol.*; 30: 201-207
24. Miller P, Osmotherly P. (2009). Does scapula taping facilitate recovery for shoulder impingement symptoms? A pilot Randomized Controlled Trial. *J Man Manipulative Ther.*; 17(1):E6-E13
25. Crossley KM, Marino GP, Macilquham MD., et al. (2009) Can patellar tape reduce the patellar malalignment and pain associated with patellofemoral osteoarthritis? *Arthritis Rheum.*; 61:1719-25
26. Lan TY, Lin WP, Jiang CC., et al. (2010). Immediate effect and predictors of effectiveness of taping for patellofemoral pain syndrome: a prospective cohort study. *Am J Sports Med.*; 38: 1626-1630
27. Gonzalez-Iglesias J, Fernandez-de-Lad-Penas C, Cleland J., et al. (2009). Short-term effects of cervical kinesio taping on pain and cervical range of motion in patients with acute whiplash injury: A randomized clinical trial. *J Orthop Sports Phys Ther.*; 39:515-521
28. Nderhand, Marc J., et al. (2002). Cervical muscle dysfunction in chronic whiplash-associated disorder grade 2: the relevance of the trauma. *Spine.* 2002; 27:1056-1061
29. Kase K. (2008). A Brief History of Kinesio Tex Taping®. Available: <http://www.kinesiotaping.co.uk/history.jsp>
30. Thelen M.D., Dauber J.A., Stoneman P.D. (2008). The Clinical Efficacy of Kinesio Tape for Shoulder Pain. *Journal of Orthopaedic and Sports Physical Therapy*; vol. 38, p. 389 – 395. ISSN 0190-6011.